



## WHAT'S NEW IN BONE AND SOFT TISSUE PATHOLOGY 2023: GUIDELINES FOR MOLECULAR TESTING

Farres Obeidin

Department of Pathology, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

Corresponding Author: Farres Obeidin, MD  
Department of Pathology, Northwestern University Feinberg School of Medicine, Chicago, IL, USA  
E-mail: farres.obeidin@nm.org

ORCID

Farres Obeidin  
<https://orcid.org/0000-0001-8461-9238>

### Abstract

Our understanding of bone and soft tissue tumors has thoroughly evolved as a consequence of modern molecular techniques. DNA and RNA sequencing methods play an important diagnostic and therapeutic role in sarcoma pathology. Herein, we discuss current guidelines and best practices for molecular testing in bone and soft tissue tumors.

### COMMON MOLECULAR METHODS

While translocation driver events are very rare in epithelial malignancies, they occur in up to 25% of sarcomas. As a result, molecular techniques for identifying recurrent translocations currently play an important diagnostic role in bone and soft tissue pathology.

### Fluorescence in-situ hybridization (FISH)

FISH uses a fluorescently labeled probe targeted towards a specific genetic sequence. The fluorescent probe can then be assessed in situ using a fluorescence microscope. Dual-color dual-fusion probes are designed towards a particular gene fusion target, while dual-color break-apart probes have greater utility for gene fusions in which the partner may not be known (e.g. *EWSR1*). Large gene amplifications and deletions may also be detected by comparing the ratio of lost target signals to a control signal (e.g. *MDM2*).

### Reverse-transcriptase polymerase chain reaction (RT-PCR)

PCR-based tests utilize PCR amplification of certain primer sequences that can be built to accommodate a set of gene rearrangements. This is particularly useful for larger panels that use known genetic breakpoints, but this method generally lacks the ability to detect new fusion partners.

### Next-generation sequencing (NGS)

NGS has become standard for detection of both prognostic and therapeutic mutations. In sarcoma, new RNA-based methods, including hybrid-capture and anchored multiplex PCR, offer better detection for gene rearrangements. These methods have the added benefit of detecting new gene partners, which is particularly useful with promiscuous genes like *EWSR1*. The development of various commercial platforms for NGS fusion testing has significantly increased the availability of this test to pathologists. Additionally, tertiary institutions with large sample volume in bone and soft tissue pathology may opt to bring these methods in-house to improve turnaround times.

### Methylation testing

Initially developed for usage in the diagnosis of brain tumors, methylation testing has expanded to begin to include soft tissue and bone tumors. Methylation tests use an array chip to detect the

methylation patterns of thousands of CpG islands to produce a “signature” for a particular tumor. Through statistical methods, these signatures can be clustered into groups of similar tumors, providing a tentative “cell of origin” diagnosis (Fig. 1). While still in its infancy, methylation has great potential for future diagnostics in soft tissue and bone.

### TESTING GUIDELINES FOR SPECIFIC TUMOR CLASSES


The constantly growing number of translocation-driven soft tissue and bone tumors in the literature necessitates judicious use of diagnostic genetic testing. By employing a morphology-based approach, pathologists can triage mesenchymal neoplasms for both diagnostic and therapeutic testing. The following review takes a “line of differentiation” approach to decide on appropriate testing strategies.

#### Adipocytic


- *MDM2* amplification is the key differentiating factor between lipoma and well-differentiated/dedifferentiated liposarcoma (Fig. 2). *MDM2* amplification may also rarely be seen in other high-grade sarcomas; as such, caution is warranted when interpreting this finding without the presence of a well-differentiated liposarcoma component.
- *RB1* deletion (tested by loss of RB1 on immunostaining or NGS) is pathognomonic for spindle cell/pleomorphic lipoma and may be seen in about 50% of atypical spindle cell/pleomorphic lipomatous tumor.
- Myxoid liposarcoma is driven by translocations involving *DDIT3*, most commonly with *FUS*. These tumors show a distinct myxoid background, chicken-wire capillary architecture, and univacuolated lipoblasts. High-grade variants show round cell features.

#### Fibroblastic/myofibroblastic

- Nodular fasciitis and similar disorganized,



**Fig. 1.** A t-distributed stochastic neighbor embedding (t-SNE) plot is a visualization method used in clinical laboratories to show the aggregate methylation data of a tumor. Each dot represents a single tumor and the closeness of the dots represents similarity in their methylation profiles. From this, a classification scheme can be created. Each cluster represents a separate tumor type. In this example case, the black dot is the current tumor and is clustering near the category of *MYOD1*-mutated rhabdomyosarcoma.



**Fig. 2.** *MDM2* amplification, seen here as an increase in red fluorescent probe signals, is a hallmark of certain cancers. The prototypical example is well-differentiated/dedifferentiated liposarcoma.

myxocollagenous, variably cellular fibroblastic/myofibroblastic neoplasms with or without reactive bone formation (including myositis ossificans, fibro-osseous pseudotumor of the digit, aneurysmal bone cyst, and a subset of fibroma of tendon sheath), form a growing spectrum of tumors that are defined by *USP6* gene rearrangements with multiple different partners (Fig. 3).

- As in spindle cell lipoma, *RB1* loss is a defining feature in both cellular angiofibroma and myofibroblastoma. These tumors have some overlapping histologic and immunophenotypic features in addition to the same genetics, so

distinction relies on a combination of clinical history and pathologic features.

- Translocation-associated fibroblastic neoplasms are typically benign to low-grade neoplasms with monotonous cytology and characteristic architectural features. An NGS panel can confirm the diagnosis, though this is often not required because the morphology is typical and molecular analogue immunohistochemical (IHC) stains exist for many of these tumors with relatively high sensitivity and specificity.
- Solitary fibrous tumor: *NAB2::STAT6* fusion; prominent collagenous background and staghorn vasculature.
- Dermatofibrosarcoma protuberans: *COL1A1::PDGFB* fusion; dermal tumor with diffuse storiform architecture and honeycomb/septal infiltration into fat. Fibrosarcomatous transformation can mask the storiform architecture and gives the tumor metastatic potential.
- Inflammatory myofibroblastic tumor: *ALK* rearrangement; bland myofibroblastic proliferation with prominent mixed inflammatory infiltrate. Malignant versions exist, often with more epithelioid morphology.
- Low-grade fibromyxoid sarcoma/sclerosing epithelioid fibrosarcoma: *FUS* or *EWSR1*

rearrangement, usually with *CREB3L2* or *CREB3L1*; cellular spindle cell neoplasm with mixed collagenous and myxoid features (low-grade fibromyxoid sarcoma) or sclerotic tumor with bland epithelioid cytology (sclerosing epithelioid fibrosarcoma). Additional fusions involving *KMT2A* and *YAP1* that show similar morphological features to sclerosing epithelioid fibrosarcoma have been found.

- *EWSR1::SMAD3*-positive fibroblastic tumor: *EWSR1::SMAD3* fusion; newly described fascicular, bland spindle cell proliferation with distinct zonation (hypocellular central region and increased peripheral cellularity) with ERG IHC positivity.
- Superficial CD34-positive fibroblastic tumor: *PRDM10* rearrangements; low-grade fascicular proliferation of spindle cells with marked nuclear pleomorphism.

### Vascular

- Fusion testing is not required but helpful in the diagnosis, particularly in the case of the low-grade malignancies and hemangioendotheliomas.
- Epithelioid hemangioma is defined by recurrent fusions involving *FOS* and *FOSB*. Fusion



**Fig. 3.** New NGS methods are highly sensitive and specific for translocations and can also be used to detect new fusions with previously undescribed partners. This example shows the readout from the NGS software demonstrating a fusion of the *USP6* gene with the *SERPINF1* gene in a case of nodular fasciitis.

testing is not required but can be useful in more cellular cases with atypical cytologic features.

- Epithelioid hemangioendothelioma: *WWTR1::CAMTA1* fusion or rarely *TFE3::YAP1* fusion; these malignant tumors show more primitive vascular differentiation, epithelioid morphology, intracytoplasmic lumens, and a distinct myxohyaline stroma that distinguishes them from epithelioid angiosarcoma. The *CAMTA1* fusion or IHC stain may be used to confirm the diagnosis.
- Pseudomyogenic hemangioendothelioma: *FOSB* gene fusions, usually with *SERPINE1* or *ACTB*; histologically, shows a rhabdomyoblastic-like appearance but stains for keratins and vascular markers.
- Angiosarcoma, like other high-grade sarcomas, shows complex genomic changes. The presence of *MYC* gene amplifications (tested by IHC, FISH, or NGS) is seen in most cases of post-irradiation or lymphedema-associated angiosarcoma. This finding is very useful in distinguishing post-irradiation atypical vascular lesions from true angiosarcoma.

### Pericytic and smooth muscle

- While the diagnosis of glomus tumors is generally based on morphology, examples of deep, gastrointestinal, or malignant glomus tumors may confound the diagnosis. Most glomus tumors (including malignant variants) show recurrent fusions involving the *NOTCH* family

of genes, most commonly *MIR143::NOTCH1/2/3*.

- Distinct from classical leiomyosarcoma, inflammatory leiomyosarcoma is a low-grade malignancy that has been shown to have a recurrent karyotypic pattern that is best seen on mRNA microarray technology, such as OncoScan™. These neoplasms show a near-haploid genotype that is thought to be a relevant driver of the disease process. Some tumors may show whole genome duplication afterwards, resulting in a pseudo-hyperdiploid karyotype that may signal transformation to a higher-grade malignancy. Recent studies have demonstrated a number of cases with rhabdomyoblastic IHC staining, and new terminology (inflammatory rhabdomyoblastic tumor) has been suggested.

### Skeletal muscle

- Molecular testing in rhabdomyoblastic tumors is best utilized in the differentiation of embryonal and alveolar rhabdomyosarcoma. Both tumors may have a solid small round blue cell morphology and skeletal muscle staining with IHC. Myogenin IHC tends to be more diffuse in alveolar rhabdomyosarcoma; however, definitive diagnosis requires molecular detection of the typical *FOXO1* fusion with either *PAX3* or *PAX7* in alveolar rhabdomyosarcoma. Embryonal rhabdomyosarcoma may show recurrent mutations in the *RAS* family of genes or *DICER1* in some syndromic patients. The distinction between the alveolar and embryonal

subtypes is necessary because of the variation in prognosis and treatment.

- Spindle cell rhabdomyosarcoma falls into three distinct molecular groupings. Congenital and infantile tumors are most often translocation-driven, with fusions involving *VGLL2*, *SRF*, *READ1*, *NCOA2*, and *CITED2*. Another subset of tumors in adolescents and young adults shows mutations in the *MYOD1* gene. The third category does not have recurrent genetic abnormalities. Additionally, some intraosseous variants may show *EWSR1*, *FUS*, or *MEIS1::NCOA2* rearrangements.

### Gastrointestinal stromal tumor (GIST)

- All GISTs should be tested for mutational status, as these mutations predict response to treatment and prognosis.
- Mutations most commonly occur in *KIT* (exons 9, 11, 13, 14, or 17) and second most commonly in *PDGFRA* (exons 12, 14, and 18).
- Immunostaining for cKIT (CD117) does not imply the presence of a *KIT* mutation.
- SDH-deficient GISTs are negative for *KIT* and *PDGFRA* mutations and show mutations in *SDHA*, *SDHB*, *SDHC*, or *SDHD*. These mutations are typically screened by assessing for loss of SDHB IHC staining, which picks up mutations in any of the four genes. SDH-deficient GISTs are more common in younger patients in the stomach.
- Some GISTs will instead show mutations in *RAS* family genes.

- A small proportion of GISTs may be negative for all four of these mutations, called quadruple wild-type GIST.

### Uncertain differentiation and round cell tumors

- Undifferentiated round to spindle cell tumors and those with monomorphic cytology should be considered for large panel NGS fusion testing. IHC has been found to show much overlap in this category of tumors, and NGS testing offers the ability to pick up non-classical examples as well as discover new fusion partners.
- Currently, Ewing and Ewing-like round cell sarcomas can be split into six overall categories (Table 1). New fusion partners continue to be discovered, and the particular fusion may affect treatment and prognosis. Certain neoplasms show bi-immunophenotypic staining patterns that may lead to confusion. Molecular testing can be confirmatory.
  - Angiomatoid fibrous histiocytoma: *EWSR1::ATF1*, *FUS::ATF1*, or *EWSR1::CREB1*; a low-grade malignancy with EMA and desmin co-positivity and prominent lymphoid cuffing.
  - Ossifying fibromyxoid tumor: Most commonly *PHF1* rearrangements (50% of cases), rarely rearrangements in *BCOR* or *SUZ12*, suggesting a genetic overlap with endometrial stromal sarcoma; low-grade malignancy with prominent peritumoral metaplastic bone formation and often co-positivity for keratins, S100, or desmin.
  - Myoepithelial neoplasms of soft tissue: *EWSR1*

or *FUS* rearrangements with several different partners; keratin and S100 co-positivity with myxoid to myxocollagenous background and bland round to spindled cells. *INI1* is lost in a subset. These tumors show different genetics to salivary gland myoepithelial neoplasms, which are often governed by *PLAG1* rearrangements.

- Extraskelatal myxoid chondrosarcoma: *NR4A3* rearrangement, most commonly with *EWSR1*. Keratin, S100, neuroendocrine, or myoepithelial markers may be nonspecifically positive. Morphologic and immunophenotypic overlap with myoepithelial neoplasms may cause diagnostic difficulty, and because of the presence of *EWSR1* as a partner in both, NGS testing is recommended to assess the partner gene to distinguish these two.
- Phosphaturic mesenchymal tumor: in patients with clinical evidence of hypophosphatemia and/or osteomalacia, serum testing may be performed for increased *FGF23* secretion. Most of these tumors show fusions involving *FN1::FGFR1* or *FN1::FGF1*.
- *NTRK* gene rearrangements have been seen in an increasing spectrum of mesenchymal tumors. The prototypical infantile fibrosarcoma is defined by *ETV6::NTRK3*. However, *NTRK* fusions have now been seen in cellular mesoblastic nephroma as well as various myxoid soft tissue tumors with bland cytology and possible co-positivity for S100 and CD34, including lipofibromatosis-like neural tumor. Some uterine and soft tissue neoplasms with fibrosarcoma-like morphology also define a new subset of *NTRK*-rearranged sarcomas.

- A subset of PEComa is driven by rearrangements in *TFE3*; often not required for the diagnosis, as the combination of myogenic and melanocytic markers is specific enough in most instances to diagnose PEComa.
- Recent studies have shown that the majority of true intimal sarcomas show amplifications in *MDM2*, similar to well/dedifferentiated liposarcoma. Intimal sarcoma may have variable differentiation and immunostaining. The presence of a luminal mass in the pulmonary or cardiac vasculature should prompt FISH testing for *MDM2* to confirm the diagnosis.

### Bone and cartilage

- The diagnosis of primary bone lesions is still based most heavily on the morphology coupled with the radiological imaging.
- Some distinct exceptions where molecular testing can be diagnostically useful:
  - Aneurysmal bone cyst: *USP6* gene rearrangements; cystic, giant cell-rich neoplasm with reactive woven bone formation.
  - Low-grade central osteosarcoma/parosteal osteosarcoma: *MDM2* amplifications; low-grade osteoblastic tumors that, similar to well-differentiated liposarcoma, have potential to dedifferentiate. *MDM2* testing by FISH can help to distinguish from reactive or benign bone-forming tumors.
  - Giant cell tumor of bone/chondroblastoma: both show unique and specific mutations in *H3F3A* or *H3F3B*. IHC testing is useful as a molecular adjunct.
  - Conventional chondrosarcoma: *IDH1* or *IDH2* mutations in a subset of chondrosarcoma; can be diagnostically useful in small biopsies and dedifferentiated examples.
  - Mesenchymal chondrosarcoma: *HEY1::NCOA2* fusion; small round blue cell sarcoma with cartilaginous maturation.

**Table 1.** Ewing and Ewing-like round cell sarcomas

Category	Molecular abnormalities
Classic Ewing sarcoma and Ewing family tumors	<i>EWSR1::FL1</i> <i>EWSR1::ERG</i> <i>EWSR1::FEV</i> <i>EWSR1::ETV1</i> <i>ESWR1::ETV4</i> <i>FUS::ERG</i> <i>FUS::FEV</i>
<i>CIC</i> -rearranged sarcoma	<i>CIC::DUX4</i> <i>CIC::FOXO4</i> <i>CIC::LEUTX</i> <i>CIC::NUTM1</i> <i>CIC::NUTM2B</i>
<i>BCOR</i> -rearranged sarcoma	<i>BCOR::CCNB3</i> <i>BCOR</i> internal tandem duplication <i>BCOR::MAML3</i>
<i>GLI1</i> -altered sarcoma	<i>GLI1::MALAT1</i> <i>GLI1::ACTB</i> <i>GLI1</i> amplifications or other rearrangements
Non-Ewing family gene fusions	<i>EWSR1::PATZ1</i> <i>EWSR1::NFATC2</i> <i>EWSR1::SP3</i> <i>EWSR1::SMARCA5</i>
Unclassified round cell sarcoma	Fusion negative or unknown

### Meet the Author

Dr. Obeidin has been an author for PathologyOutlines since 2018 and part of the PathologyOutlines editorial board since January 2022. He is currently an Assistant Professor of Pathology at Northwestern University Feinberg School of Medicine. He obtained his M.D. at the Medical College of Georgia and then completed his Anatomic and Clinical Pathology residency at Northwestern University. He then completed a fellowship in General Surgical Pathology and Bone and Soft Tissue Pathology at the University of California, Los Angeles.